

3B

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 31/64, 31/70, 31/715 // (A61K 31/64, 31:715, 31:70)</b>		<b>A1</b>	(11) International Publication Number: <b>WO 99/03478</b>
			(43) International Publication Date: 28 January 1999 (28.01.99)
(21) International Application Number: <b>PCT/GB98/02112</b>		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 16 July 1998 (16.07.98)		<p><b>Published</b> With international search report.</p>	
(30) Priority Data: 9715298.7 18 July 1997 (18.07.97) GB			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BUCKINGHAM, Robin, Edwin [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). SMITH, Stephen, Alistair [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).			
(74) Agent: RUTTER, Keith; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(54) Title: TREATMENT OF DIABETES WITH THIAZOLIDINEDIONE, INSULIN SECRETAGOGUE AND ALPHA GLUCOCIDASE INHIBITOR			
(57) Abstract			
<p>A method for the treatment of diabetes mellitus and conditions associated with diabetes mellitus in a mammal, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent, to a mammal in need thereof, and composition for use in such method.</p>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**TREATMENT OF DIABETES WITH THIAZOLIDINEDIONE, INSULIN SECRETAGOGUE AND ALPHA GLUCOCIDASE INHIBITOR**

This invention relates to a method of treatment, in particular to a method for the treatment of diabetes mellitus, especially non-insulin dependent diabetes  
5 (NIDDM) or Type 2 diabetes and conditions associated with diabetes mellitus.

Insulin secretagogues are compounds which promote increased secretion of insulin by the pancreatic beta cells.

The sulphonylureas are well known examples of insulin secretagogues. The sulphonylureas act as antihyperglycaemic agents and are used in the treatment of Type  
10 2 diabetes. Examples of sulphonylureas include glibenclamide, glipizide, gliclazide, glimepiride, tolazamide and tolbutamide.

Alpha glucosidase inhibitor antihyperglycaemic agents, such as acarbose, emiglitate and miglitol, are commonly used in the treatment of Type 2 diabetes.

European Patent Application. Publication Number 0,306,228 relates to certain  
15 thiazolidinedione derivatives disclosed as having antihyperglycaemic and antihyperlipidaemic activity. One particular thiazolidinedione disclosed in EP 0306228 is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (hereinafter 'Compound (I)'). WO94/05659 discloses certain salts of Compound (I) including the maleate salt.

20 Compound (I) is an example of a class of anti-hyperglycaemic agents known as 'insulin sensitisers'. In particular Compound (I) is a thiazolidinedione insulin sensitiser.

European Patent Applications, Publication Numbers: 0008203, 0139421, 0032128, 0428312, 0489663, 0155845, 0257781, 0208420, 0177353, 0319189,  
25 0332331, 0332332, 0528734, 0508740; International Patent Application, Publication Numbers 92/18501, 93/02079, 93/22445 and United States Patent Numbers 5104888 and 5478852, also disclose certain thiazolidinedione insulin sensitisers.

Another series of compounds generally recognised as having insulin sensitiser activity are those typified by the compounds disclosed in International Patent  
30 Applications, Publication Numbers WO93/21166 and WO94/01420. These compounds are herein referred to as 'acyclic insulin sensitisers'. Other examples of acyclic insulin sensitisers are those disclosed in United States Patent Number 5232945 and International Patent Applications. Publication Numbers WO92/03425 and WO91/19702.

35 Examples of other insulin sensitisers are those disclosed in European Patent Application. Publication Number 0533933, Japanese Patent Application Publication Number 05271204 and United States Patent Number 5264451.

The above mentioned publications are incorporated herein by reference.

provides a particularly beneficial effect on glycaemic control, such combination is therefore particularly useful for the treatment of diabetes mellitus, especially Type 2 diabetes, and conditions associated with diabetes mellitus. The treatment is also indicated to proceed with minimum side effects.

Accordingly, the invention provides a method for the treatment of diabetes mellitus, especially Type 2 diabetes, and conditions associated with diabetes mellitus, in a mammal such as a human, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent, to a mammal in need thereof.

The method comprises either co-administration of the insulin sensitiser, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent or sequential administration thereof.

Co-administration includes administration of a formulation which includes an insulin sensitiser, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent or the essentially simultaneous administration of separate formulations of each agent.

In another aspect the invention provides the use of an insulin sensitiser, such as Compound (I), an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent, in the manufacture of a composition for the treatment of diabetes mellitus, especially Type 2 diabetes and conditions associated with diabetes mellitus.

A suitable insulin sensitiser is a thiazolidinedione insulin sensitiser.

A suitable thiazolidinedione insulin sensitiser is Compound (I).

Other suitable thiazolidinedione insulin sensitisers include (+) -5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl)methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione (or pioglitazone) or 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone)

Suitable insulin secretagogues include sulphonylureas.

Suitable sulphonylureas include glibenclamide, glipizide, gliclazide, glimepiride, tolazamide and tolbutamide.

Further sulphonylureas include acetohexamide, carbutamide, chlorpropamide, glibornuride, gliquidone, glisentide, glisolamide, glisoxepide, glyclopyamide and glycylamide.

Further suitable insulin secretagogues include repaglinide.

A suitable alpha glucosidase inhibitor antihyperglycaemic agent is acarbose.

Other suitable alpha glucosidase inhibitor antihyperglycaemic agents are emiglitate and miglitol.

- 5 In one particular aspect, the method comprises the administration of 2 to 12 mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 2 to 4, 4 to 8 or 8 to 12 mg of Compound (I) per day.

- 10 Particularly, the method comprises the administration of 2 to 4mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 4 to 8mg of Compound (I), especially when administered per day.

Particularly, the method comprises the administration of 8 to 12 mg of Compound (I), especially when administered per day.

- 15 ~~Preferably, the method comprises the administration of 2 mg of Compound (I), especially when administered per day.~~

Preferably, the method comprises the administration of 4 mg of Compound (I), especially when administered per day.

- 20 Preferably, the method comprises the administration of 8 mg of Compound (I), especially when administered per day.

It will be understood that the insulin sensitiser, such as compound (I), the insulin secretagogue and the alpha glucosidase inhibitor antihyperglycaemic agent are each administered in a pharmaceutically acceptable form, including pharmaceutically acceptable derivatives such as pharmaceutically acceptable salts, esters and solvates thereof, as appropriate. In certain instances herein the names used for the relevant  
25 insulin secretagogues and the alpha glucosidase inhibitor antihyperglycaemic agents may relate to a particular pharmaceutical form of the relevant active agent: It will be understood that all pharmaceutically acceptable forms of the active agents per se are encompassed by this invention.

- 30 Suitable pharmaceutically acceptable salted forms of the insulin sensitisers, such as Compound (I), include those described in the above mentioned patents and patent applications such as in EP 0306228 and WO94/05659 for Compound (I). A preferred pharmaceutically acceptable salt for Compound (I) is a maleate.

- 35 Suitable pharmaceutically acceptable solvated forms of the insulin sensitisers, such as Compound (I), include those described in the above mentioned patents and patent applications, such as in EP 0306228 and WO94/05659 for Compound (I), in particular hydrates.

Suitable pharmaceutically acceptable forms of the insulin secretagogue and the alpha glucosidase inhibitor antihyperglycaemic agent depend upon the particular compound used but include known pharmaceutically acceptable forms of the particular compound chosen. Such derivatives are found or are referred to in standard  
5 reference texts such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages cited therein).

The insulin sensitisers, such as Compound (I) or, a pharmaceutically  
10 acceptable salt thereof, or a pharmaceutically acceptable solvate thereof, may be prepared using known methods, for example those disclosed in the above mentioned patents and patent applications, such as EP 0306228 and WO94/05659 for Compound (I). The disclosures of the above mentioned patents and patent applications, such as EP 0306228 and WO94/05659, are incorporated herein by reference.

15 Compound (I) may exist in one of several tautomeric forms, all of which are encompassed by the term Compound (I) as individual tautomeric forms or as mixtures thereof. Compound (I) contains a chiral carbon atom, and hence can exist in up to two stereoisomeric forms, the term Compound (I) encompasses all of these isomeric forms whether as individual isomers or as mixtures of isomers, including racemates.

20 The insulin secretagogue and alpha glucosidase inhibitor antihyperglycaemic agent of choice is prepared according to known methods, such methods are found or are referred to in standard reference texts, such as the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition  
25 page 341 and pages cited therein).

When used herein the term 'conditions associated with diabetes' includes conditions associated with diabetes mellitus itself and complications associated with diabetes mellitus. Also included in 'conditions associated with diabetes' are those conditions associated with the pre-diabetic state.

30 When used herein the term 'conditions associated with the pre-diabetic state' includes conditions such as insulin resistance, including hereditary insulin resistance, impaired glucose tolerance and hyperinsulinaemia.

'Conditions associated with diabetes mellitus itself' include hyperglycaemia, insulin resistance, including acquired insulin resistance. Further conditions associated  
35 with diabetes mellitus itself include hypertension and cardiovascular disease, especially atherosclerosis and conditions associated with insulin resistance. Conditions associated with insulin resistance include polycystic ovarian syndrome and steroid induced insulin resistance and gestational diabetes.

'Complications associated with diabetes mellitus' includes renal disease, especially renal disease associated with Type 2 diabetes, neuropathy and retinopathy.

Renal diseases associated with Type 2 diabetes include nephropathy, glomerulonephritis, glomerular sclerosis, hypertensive nephrosclerosis and end stage renal disease. Additional renal diseases associated with Type 2 diabetes include nephrotic syndrome.

For the avoidance of doubt, when reference is made herein to scalar amounts, including mg amounts, of Compound (I) in a pharmaceutically acceptable form, the scalar amount referred to is made in respect of Compound (I) *per se*: For example 2 mg of Compound (I) in the form of the maleate salt is that amount of maleate salt which contains 2 mg of Compound (I).

Diabetes mellitus is preferably Type 2 diabetes.

The particularly beneficial effect on glycaemic control provided by the treatment of the invention is indicated to be a synergistic effect relative to the control expected for the sum of the effects of the individual active agents.

Glycaemic control may be characterised using conventional methods, for example by measurement of a typically used index of glycaemic control such as fasting plasma glucose or glycosylated haemoglobin (HbA1c). Such indices are determined using standard methodology, for example those described in: Tiescher A, Richterich, P., Schweiz. med. Wschr. 101 (1971), 345 and 390 and Frank P., 'Monitoring the Diabetic Patient with Glycosylated Hemoglobin Measurements', Clinical Products 1988

In a preferred aspect, the dosage level of each of the active agents when used in accordance with the treatment of the invention will be less than would have been required from a purely additive effect upon glycaemic control.

Suitably, the insulin sensitiser is the agent of first administration.

Suitably the insulin secretagogue is the agent of second administration.

Suitably the alpha glucosidase inhibitor is the agent of third administration.

There is also an indication that the treatment of the invention will effect an improvement, relative to the individual agents, in the levels of advanced glycosylation end products (AGEs), leptin and serum lipids including total cholesterol, HDL-cholesterol, LDL-cholesterol including improvements in the ratios thereof, in particular an improvement in serum lipids including total cholesterol, HDL-cholesterol, LDL-cholesterol including improvements in the ratios thereof.

As used herein the term 'pharmaceutically acceptable' embraces both human and veterinary use: for example the term 'pharmaceutically acceptable' embraces a veterinarily acceptable compound.

In the method of the invention, the active medicaments are preferably administered in pharmaceutical composition form. As indicated above, such compositions can include all medicaments or one only of the medicaments.

Accordingly, in one aspect the present invention also provides a  
5 pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.

Such compositions may be prepared by admixing an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and an  
10 alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.

Usually the compositions are adapted for oral administration. However, they may be adapted for other modes of administration, for example parenteral administration, sublingual or transdermal administration.

15 The compositions may be in the form of tablets, capsules, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid preparations, such as oral or sterile parenteral solutions or suspensions.

In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit dose.

20 Unit dose presentation forms for oral administration may be tablets and capsules and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for example magnesium stearate; disintegrants, for example  
25 starch, polyvinylpyrrolidone, sodium starch glycollate or microcrystalline cellulose; or pharmaceutically acceptable wetting agents such as sodium lauryl sulphate.

The compositions are preferably in a unit dosage form in an amount appropriate for the relevant daily dosage.

Suitable dosages for the insulin sensitisers include those disclosed in the  
30 abovementioned patents and patent applications.

Suitable dosages, including unit dosages, of Compound (I) comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 mg of Compound (I).

In the treatment the medicaments may be administered from 1 to 6 times a day, but most preferably 1 or 2 times per day.

35 Particular dosages of Compound (I) are 2mg/day, 4mg/day, including 2mg twice per day, and 8 mg/day, including 4mg twice per day.

Suitable dosages including unit dosages of the insulin secretagogue, such as the sulphonylurea, or the alpha glucosidase inhibitor antihyperglycaemic agent,



include the known dosages including unit doses for these compounds as described or referred to in reference text such as the British and US Pharmacopoeias. Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) (for example see the 31st Edition page 341 and pages cited therein).

Thus for the sulphonylureas, a typical daily dosage of glibenclamide is in the range of from 2.5 to 20 mg, for example 10mg twice per day or 20mg once per day; a typical daily dosage of glipizide is in the range of from 2.5 to 40 mg; a typical daily dosage of gliclazide is in the range of from 40 to 320 mg; a typical daily dosage of tolazamide is in the range of from 100 to 1000 mg; a typical daily dosage of tolbutamide is in the range of from 1000 to 3000 mg; a typical daily dosage of chlorpropamide is in the range of from 100 to 500 mg; and a typical daily dosage of gliquidone is in the range of from 15 to 180 mg.

Repaglinide may be taken in amounts, usually in the range of from 0.5mg to 4mg and usually with meals, up to a typical maximum daily dosage of 16mg per day.

With regard to the alpha glucosidase inhibitor antihyperglycaemic agents, a typical daily dosage of acarbose is in the range of from 50 to 600 mg, an example 100mg or 200mg per day.

The solid oral compositions may be prepared by conventional methods of blending, filling or tableting. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are of course conventional in the art. The tablets may be coated according to methods well known in normal pharmaceutical practice, in particular with an enteric coating.

Oral liquid preparations may be in the form of, for example, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethylcellulose, aluminium stearate gel, hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid; and if desired conventional flavouring or colouring agents.

For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, and, depending on the concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the compound can

be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, a preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the Compound (I) is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration. The compound can be sterilized by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

Compositions may contain from 0.1% to 99% by weight, preferably from 10-60% by weight, of the active material, depending upon the method of administration.

Composition may, if desired, be in the form of a pack accompanied by written or printed instructions for use.

The compositions are prepared and formulated according to conventional methods, such as those disclosed in standard reference texts, for example the British and US Pharmacopoeias, 's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Extra Pharmacopoeia (London, The Pharmaceutical Press) and Harry's Cosmeticology (Leonard Hill Books) (for example see the 31st Edition page 341 and pages cited therein).

The present invention also provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic substance.

The invention also provides the use of an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent for the manufacture of a medicament for the treatment of diabetes mellitus and conditions associated with diabetes mellitus.

In particular, the present invention provides a pharmaceutical composition comprising an insulin sensitiser, such as Compound (I) and especially 2 to 12 mg thereof, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use in the treatment of diabetes mellitus and conditions associated with diabetes mellitus.

A range of 2 to 4mg includes a range of 2.1 to 4, 2.2 to 4, 2.3 to 4, 2.4 to 4, 2.5 to 4, 2.6 to 4, 2.7 to 4, 2.8 to 4, 2.9 to 4 or 3 to 4mg.

A range of 4 to 8mg includes a range of 4.1 to 8, 4.2 to 8, 4.3 to 8, 4.4 to 8, 4.5 to 8, 4.6 to 8, 4.7 to 8, 4.8 to 8, 4.9 to 8, 5 to 8, 6 to 8 or 7 to 8mg.

A range of 8 to 12 mg includes a range of 8.1 to 12, 8.2 to 12, 8.3 to 12, 8.4 to 12, 8.5 to 12, 8.6 to 12, 8.7 to 12, 8.8 to 12, 8.9 to 12, 9 to 12, 10 to 12 or 11 to 12mg.

No adverse toxicological effects are expected for the compositions or methods  
5 of the invention in the abovementioned dosage ranges.

**Composition of Compound (I)**

**Preparation of Concentrate:** Tableting concentrate was prepared using the following materials

Ingredient	Quantity (%)
Milled Compound (I) as maleate salt	13.25 (pure maleate salt)
Sodium Starch Glycollate	5.00
Hydroxypropyl Methylcellulose 2910	5.00
Microcrystalline Cellulose (Avicel PH102)	20.0
Lactose Monohydrate, regular grade	to 100
Purified water	*

\* Removed during processing.

The concentrate was then formulated into tablets using the following:

Tablet Strength	Quantity (mg per Tablet)			
	1.0mg	2.0mg	4.0mg	8.0mg
<b>Active Ingredient:</b>				
Compound (I) maleate Concentrate granules	10.00	20.00	40.00	80.00
<b>Other Ingredients:</b>				
Sodium Starch Glycolate	6.96	6.46	5.46	10.92
Microcrystalline Cellulose (Avicel PH102)	27.85	25.85	21.85	43.70
Lactose monohydrate, (Pharmatose DCL15),	104.44	96.94	81.94	163.88
Magnesium Stearate	0.75	0.75	0.75	1.50
<b>Total Weight of Tablet Core</b>	<b>150.0</b>	<b>150.0</b>	<b>150.0</b>	<b>300.0</b>
Opadry	4.5	4.5	4.5	9.0
<b>Total Weight of Film Coated Tablet</b>	<b>154.5</b>	<b>154.5</b>	<b>154.5</b>	<b>309.0</b>

Compositions of the other active agents are as disclosed in the references mentioned herein.

## Claims

1. A method for the treatment of diabetes mellitus and conditions associated with diabetes mellitus in a mammal, which method comprises administering an effective non-toxic and pharmaceutically acceptable amount of an insulin sensitiser, an insulin secretagogue and an alpha glucosidase inhibitor antihyperglycaemic agent, to a mammal in need thereof.
2. A method according to claim 1, wherein the insulin secretagogue is glibenclamide, glipizide, gliclazide, glimepiride, tolazamide, tolbutamide, acetohexamide, carbutamide, chlorpropamide, glibornuride, gliquidone, glisentide, glisolamide, glisoxepide, glyclopyamide, glycylamide or repaglinide.
3. A method according to claim 1, wherein the alpha glucosidase inhibitor antihyperglycaemic agent is acarbose, emiglitate and miglitol.
4. A method according to any one of claims 1 to 3, wherein the insulin sensitiser is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (Compound I).
5. A method according to any one of claims 1 to 4, which comprises the administration of 2 to 12 mg of Compound (I).
6. A method according to any one of claims 1 to 5, which comprises the administration of 2 to 4, 4 to 8 or 8 to 12 mg of Compound (I).
7. A method according to any one of claims 1 to 6, which comprises the administration of 2 to 4mg of Compound (I).
8. A method according to any one of claims 1 to 6, which comprises the method the administration of 4 to 8mg of Compound (I).
9. A method according to any one of claims 1 to 6, which comprises the administration of 8 to 12 mg of Compound (I).
10. A method according to any one of claims 1 to 6, which comprises the administration of 2 mg of Compound (I).

11. A method according to any one of claims 1 to 6, which comprises the administration of 4 mg of Compound (I).
12. A method according to any one of claims 1 to 6, which comprises the administration of 8 mg of Compound (I).
13. A method according to claim 1, wherein the insulin sensitiser is (+) -5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl]methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione (or pioglitazone) or 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone); or a pharmaceutically acceptable form thereof.
14. A pharmaceutical composition comprising an insulin sensitiser, an insulin secretagogue, an alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor.
15. A composition according to claim 14, wherein the insulin secretagogue is a sulphonylurea.
16. A composition according to claim 14 or claim 15, wherein the insulin secretagogue is glibenclamide, glipizide, gliclazide, glimepiride, tolazamide or tolbutamide, acetohexamide, carbutamide, chlorpropamide, glibornuride, gliquidone, glisentide, glisolamide, glisoxepide, glycopyramide, glycylamide or repaglinide.
17. A composition according to claim 1, wherein the alpha glucosidase inhibitor antihyperglycaemic agent is acarbose, emiglitate and miglitol.
18. A composition according to any one of claims 14 to 17, wherein the insulin sensitiser is Compound (I)
19. A composition according to any one of claims 14 to 17, which comprises 2 to 12 mg of Compound (I).

20. A pharmaceutical composition comprising an insulin sensitiser an insulin secretagogue and a pharmaceutically acceptable carrier therefor, for use as an active therapeutic substance.
- 5 21. A pharmaceutical composition comprising an insulin sensitiser, an insulin secretagogue, an alpha glucosidase inhibitor antihyperglycaemic agent and a pharmaceutically acceptable carrier therefor, for use in the treatment of diabetes mellitus and conditions associated with diabetes mellitus.
- 10 22. A composition according to any one of claims 14, 20 or 21, wherein the insulin sensitiser is (+) -5-[[4-[(3,4-dihydro-6-hydroxy-2, 5, 7, 8-tetramethyl-2H-1-benzopyran-2-yl)methoxy]phenyl)methyl]-2,4-thiazolidinedione (or troglitazone), 5-[4-[(1-methylcyclohexyl)methoxy]benzyl] thiazolidine-2,4-dione (or ciglitazone), 5-[4-[2-(5-ethylpyridin-2-yl)ethoxy]benzyl] thiazolidine-2,4-dione
- 15 ~~(or pioglitazone) or 5-[(2-benzyl-2,3-dihydrobenzopyran)-5-ylmethyl]thiazolidine-2,4-dione (or englitazone);~~ or a pharmaceutically acceptable form thereof.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/02112

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/64 A61K31/70 A61K31/715 //(A61K31/64.31:715.31:70)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
Y	EP 0 749 751 A (TAKEDA CHEMICAL INDUSTRIES LTD) 27 December 1996 see page 9, line 42-55; claims 24-26; example 2 see page 8, line 32-34	1-22
Y,P	KAPPEL C. ET AL: "Type 2 diabetes: Update on therapy" COMPREHENSIVE THERAPY. vol. 24, no. 6-7, June 1998, pages 319-326, XP002080974 see page 324, column 2, paragraph 2; figure 1	1-22

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 October 1998

Date of mailing of the international search report

05/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gonzalez Ramon, N

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/02112

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	SCHEEN AJ ET AL: "Oral antidiabetic agents. A guide to selection." DRUGS, FEB 1998, 55 (2) P225-36, XP002080821 NEW ZEALAND see page 232, column 2, paragraph 1; figure 1 ---	1-22
A	WO 93 03724 A (UPJOHN CO) 4 March 1993 see claims 3,4,10,11 ---	1-22
A	WO 96 09823 A (SHAMAN PHARMACEUTICALS INC) 4 April 1996 see claims, 7-11 ---	1-22
A	ELSON DF ET AL: "Therapy for type 2 diabetes mellitus 'see comments!'" WMJ, MAR 1998, 97 (3) P49-54, XP002081124 UNITED STATES see page 52	1-22
Y	see page 53, paragraph 2 ---	1-22
A	WOLFENBUTTEL B. ET AL: "New treatments for patients with type 2 diabetes mellitus" POSTGRAD MED, vol. 72, 1996, pages 657-662, XP002081125 boxes 3,4,5 see abstract -----	1-22

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB 98/02112

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos. because they relate to subject matter not required to be searched by this Authority namely:  
 Remark: Although claims 1-13 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos. because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos. because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Inte. Appl. Application No

PCT/GB 98/02112

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0749751 A	27-12-1996	AU 5603496 A	09-01-1997
		CA 2179584 A	21-12-1996
		CN 1145783 A	26-03-1997
		CZ 9601811 A	15-01-1997
		EP 0861666 A	02-09-1998
		HU 9601698 A	28-05-1997
		JP 9067271 A	11-03-1997
		JP 10167986 A	23-06-1998
		NO 962606 A	23-12-1996
		SK 79496 A	08-01-1997
WO 9303724 A	04-03-1993	AT 151634 T	15-05-1997
		AU 2479492 A	16-03-1993
		DE 69219136 D	22-05-1997
		DE 69219136 T	28-08-1997
		DK 601001 T	20-10-1997
		EP 0601001 A	15-06-1994
		ES 2101866 T	16-07-1997
		GR 3023848 T	30-09-1997
		JP 6510286 T	17-11-1994
WO 9609823 A	04-04-1996	AU 3731895 A	19-04-1996
		AU 3731995 A	19-04-1996
		WO 9610015 A	04-04-1996
		US 5628999 A	13-05-1997
		US 5753790 A	19-05-1998
		US 5629319 A	13-05-1997